

# Geometry

## 4.1 Translations


### Vector ( $\vec{BC}$ )

Measurement with \_\_\_\_\_ and \_\_\_\_\_ (size)




Represented by an \_\_\_\_\_

Component form  $\langle$  \_\_\_\_\_, \_\_\_\_\_  $\rangle$

$\vec{BC} = \langle$  \_\_\_\_\_, \_\_\_\_\_  $\rangle$



Name the vector and write its component form



### Transformation

\_\_\_\_\_ or \_\_\_\_\_ a figure

Original called \_\_\_\_\_ (i.e.  $\triangle ABC$ )

New called \_\_\_\_\_ (i.e.  $\triangle A'B'C'$ )

### Translation


\_\_\_\_\_ every point the same \_\_\_\_\_ in the same \_\_\_\_\_

$(x, y) \rightarrow$  \_\_\_\_\_

Where \_\_\_\_\_ is the translation \_\_\_\_\_


The vertices of  $\triangle LMN$  are  $L(2, 2)$ ,  $M(5, 3)$ ,  $N(9, 1)$ . Translate  $\triangle LMN$  using vector  $\langle -2, 6 \rangle$ .

Write a rule for the translation of  $\triangle PQR$  to  $\triangle P'Q'R'$ .



Draw  $\triangle RST$  with vertices  $R(2, 2)$ ,  $S(5, 2)$ , and  $T(3, -2)$ . Find the image of each vertex after the translation  $(x, y) \rightarrow (x + 1, y + 2)$ .

Graph the image using prime notation.



**Rigid Motion**

Transformation that preserves \_\_\_\_\_ and \_\_\_\_\_.

A \_\_\_\_\_ transformation

**Translation Theorem**  
A translation is a \_\_\_\_\_.

**Composition of Transformations**

- \_\_\_\_\_ or \_\_\_\_\_ transformations \_\_\_\_\_ into a \_\_\_\_\_ transformation

**Composition Theorem**  
A composition of two (or more) \_\_\_\_\_ is a \_\_\_\_\_.

Graph  $\overline{RS}$  with endpoints  $R(-8, 5)$  and  $S(-6, 8)$ . Graph its image after the composition.

**Translation:**  $(x, y) \rightarrow (x - 1, y + 4)$

**Translation:**  $(x, y) \rightarrow (x + 4, y - 6)$

