

Clements 2004 World Taxonomy Changes

July 15, 2003. Page 19

Race *gularis* of the Little Egret is raised to species status.

Western Reef-Heron *Egretta gularis*. Range of nominate race and *schistacea* as published in the checklist. Cramp, Stanley et al. editors. 1977. Handbook of the Birds of Europe, the Middle East and North Africa. The Birds of the Western Palearctic. Volume 1: Ostrich to Ducks. Oxford University Press.

July 15, 2003. Page 49

After Lined Forest-Falcon add Cryptic Forest-Falcon *Micrastur mintoni*. Range: Rainforests of Brazil and adjacent northeast Bolivia. Whittaker, Andrew. 2002. A new species of forest-falcon (Falconidae: *Micrastur*) from southeastern Amazonia and the Atlantic rainforests of Brazil. *Wilson Bulletin* 114 (4): 421-445.

July 15, 2003. Page 60

Under Greater Prairie-Chicken change range of race *T. c. pinnatus* to read: South-central Canada to northeast Oklahoma and northern Texas. American Ornithologists Union. 1998. Check-list of North American Birds. Seventh Edition.

July 15, 2003. Page 68

Under Udzungwa Partridge add race *X. u. obscurata*. Range: N Tanzania (Maf-wemiro forest in Rubeho highlands). Fjeldså, Jon and Jacob Kiure. 2003. A new population of the Udzungwa Forest Partridge. *Bull. B. O. C.* 123 (1): 52-57.

July 15, 2003. Page 167

Vermiculated Screech-Owl *Otus vermiculatus* now considered monotypic. Range: Lowlands of Costa Rica to new Colombia and n Venezuela. Race *roraimae* raised to species level. *Roraima Screech-Owl Otus roraimae*. Monotypic. Range: Tepuis of se Venezuela and adjacent n Brazil. Race *napensis* raised to species level. *Rio Napo Screech-Owl Otus napensis* with following races. *O. n. napensis*, range: Tropical e Ecuador and e Colombia; *O. n. helleri*, range: Tropical e Peru; *O. n. bolivianus*, range: Tropical n Bolivia (Cochabamba). König, Claus, Friedhelm Weick and Jan-Hendrik Becking. 1999. *Owls: A Guide to the Owls of the World*. Sussex, UK: Pica Press.

July 15, 2003. Page 173

After Least Pygmy-Owl add Pernambuco Pygmy-Owl *Glaucidium mooreorum*. Range: Known only from two locations in southern Pernambuco, eastern Brazil. Galileu, Arthur and Luiz Gonzaga. 2003. A New Pygmy-Owl from Pernambuco. Ararajuba. In press.

July 15, 2003. Page 176

The race japonica of the Brown Hawk-Owl is raised to species level. Northern Boobook *Ninox japonica*. Range: SE Siberia to se Manchuria, e China, Korea and Japan. Add race *Ninox j. totogo* with range: Ryukyu Islands and Taiwan. Ben King suggests that totogo might be specifically distinct. Race *ussuriensis* considered synonymous with nominate race. Race *randi* is raised to species level. Chocolate Boobook *Ninox randi*. Monotypic. Range: Philippine Islands (except Palawan), Sulu Archipelago and Talaud Archipelago. King, Ben. 2002. Species limits in the Brown Boobook *Ninox scutulata* complex. Bull. B. O. C. 122 (4): 250-257.

July 15, 2003. Page 244

Correct spelling of *Eubucco richardsoni aurantiicollis*. Change range of *Eubucco bourcierii aequatorialis* to: Central and E Andes of Colombia and Andes of w Venezuela. Change range of *Eubucco bourcierii orientalis* to read: Subtropical e slope of Andes of Ecuador and n Peru.

July 15, 2003. Page 245

The Emerald Toucanet is split into seven species, with ranges as follows:

Wagler's Toucanet *Aulacorhynchus wagleri*. Monotypic. Range: Sierra Madre del Sur of sw Mexico (Guerrero and sw Oaxaca)

Emerald Toucanet *Aulacorhynchus prasinus*

A. p. prasinus Range: SE Mexico (Veracruz, adj. San Luis Potosí and Oaxaca)

A. p. warneri Range: Mts. of se Mexico (Sierra de Los Tuxtlas in s Veracruz)

A. p. virescens Range: SE Mexico (Chiapas) to Honduras and Nicaragua

A. p. stenorhabdus Range: Subtropical s Mexico to w Guatemala and n El Salvador

A. p. chiapensis Range: Mts. of extreme s Mexico (Mt. Ovando, Chiapas)

A. p. volcanius Range: Eastern El Salvador (Volcán San Miguel)

Blue-throated Toucanet *Aulacorhynchus caeruleogularis*

A. c. maxillaris Range: Highlands of Costa Rica and w Panama

A. c. caeruleogularis Range: Highlands of e-central Panama (Chiriquí and Veraguas)

Violet-throated Toucanet *Aulacorhynchus cognatus*. Monotypic. Range: Mts. of e Panama (Darién) and adjacent Colombia

Santa Marta Toucanet *Aulacorhynchus lautus*. Monotypic. Range: Santa Marta Mountains (ne Colombia)

Andean Toucanet *Aulacorhynchus albivitta*

A. a. albivitta Range: E and Central Andes of Colombia, e Ecuador and w Venezuela

A. a. griseigularis Range: N end of W Andes and w slope of Central Andes of Colombia

A. a. phaeolaemus (petax) Range: Subtropical W Andes of Colombia

A. a. cyanolaemus Range: Subtropical Andes of se Ecuador and n Peru

Black-throated Toucanet *Aulacorhynchus atrogularis*

A. a. atrogularis Range: Humid Andes of e Peru and n Bolivia

A. a. dimidiatus Range: Subtropical Andes of n Peru

Navarro, Adolfo G., A. Townsend Peterson, Esteban López-Medrano and Hesequio Benítez-Díaz. 2001. Species limits in Mesoamerican *Aulacorhynchus* Toucanets. *Wilson Bulletin* 114 (4): 363-372.

July 15, 2003. Page 301

Parana Antwren is moved after Narrow-billed Antwren, and placed in genus *Formicivora*. Gonzaga, L. P. 2001. Análise filogenética do Género *Formicivora* Swainson 1825 (Aves: Passeriformes, *Thamnophilidae*) baseada em caracteres morfológicos e vocais. *Resumo de tese. Atualidades Ornitológicas* 102: 2.

Narrow-billed Antwren is placed in a monotypic genus *Neorhophias* and follows Dot-winged Antwren in taxonomic order. *Neorhophias iheringi*.

July 15, 2003. Page 374

After Malindi Pipit add Kimberley Pipit *Anthus pseudosimilis*. Range: Kimberley and adjacent regions of South Africa. Liversidge, Richard and Gary Voekler. 2002. The Kimberley Pipit: a new African species. *Bull. B. O. C.* 122 (2): 93-109.

July 15, 2003. Page 390

Race *notata* with subspecies *ugandae* of Green-tailed Bristlebill is raised to species level. Lesser Bristlebill *Bleda notata*. Ranges as published in the checklist. Borrow, Nik and Ron Demey. 2001. *A Guide to the Birds of Western Africa*. Princeton University Press.

July 15, 2003. Page 400

Under Bay Wren, add race *T. n. odicus* after *T. n. castaneus*. Range: Panama (Escudo de Veraguas island). Brewer, David. 2001. *Wrens, Dippers and Thrashers*. Yale University Press.

July 15, 2003. Page 404

Race *festinus* of Mountain Wren is now considered a race of Ochraceous Wren *Troglodytes o. festinus*. Range: E Panama (Mt. Pirre) and possibly adjacent Colombia. Ridgely, Robert S. 1976. *A Guide to the Birds of Panama*. Princeton University Press.

July 15, 2003. Page 406

Under Scaly-breasted Wren, changes races as follows: *M. m. luscini*a: Central Costa Rica to e Panama and extreme nw Colombia; *M. m. taeniatus*: Tropical w Ecuador; *M. m. marginatus*: E Colombia to n Bolivia and w Amazonian Brazil; *M. m. squamulatus*: NE Colombia and nw Venezuela; *M. m. corrasus*: Santa Marta region of n Colombia; *M. m. occidentalis*: W Colombia and nw Ecuador. Brewer, David. 2001. Wrens, Dippers and Thrashers. Yale University Press and Ridgely, Robert S. and Paul J. Greenfield. 2001. The Birds of Ecuador: Status, Distribution and Taxonomy. Cornell University Press.

July 15, 2003. Page 447

Under Iberian Chiffchaff add following subspecies: *P. i. biscayensis*, Range: Northern Portugal and n Spain to Basque France. *P. i. ibericus*, Range: Central and s Portugal to sw Spain (Andalucia). Race *tristis* becomes a race of *P. collybita* with range as published in the checklist. Salomon, Marc, Jean-François Voisin and Joël Bried. 2002. On the taxonomic status and denomination of the Iberian Chiffchaffs. Ibis 145: 87-97.

Change scientific name of Iberian Chiffchaff to *Phylloscopus ibericus*. Svensson, Lars. 2001. The correct name of the Iberian Chiffchaff *Phylloscopus ibericus* Ticehurst 1937, its identification and new evidence of its winter grounds. Bulletin B. O. C. 121 (4): 281-296.

July 15, 2003. Page 448

Under Arctic Warbler correct spelling of *P. b. kennicotti*. American Ornithologists Union. 1957. Check-list of North American Birds. Fifth Edition.

July 15, 2003. Page 472

Change English name of Canary Islands Chat to Canary Island Stonechat as the word "Stonechat" is specific to the genus. The species is now confined to a single island in the Canaries (Fuerteventura) thus "Islands" is inappropriate. Urquhart, Ewan. 2002. Stonechats: A Guide to the Genus *Saxicola*. Yale University Press.

The English name of *Saxicola rubicola* is changed to European Stonechat, with races *rubicola* and *hibernans*. Ranges as published in the checklist. Ibid.

After European Stonechat add Siberian Stonechat *Saxicola maura* with races *variegata*, *armenica*, *indica*, *stejnegeri*, and *przewalskii*. Ibid.

Add following races to African Stonechat. *S. t. adamauae*, Range:

Highlands of n and w Cameroon; and *S. t. pallidigula* Range: Cameroon Mt. and Bioko. Ibid.

July 15, 2003. Page 473

Change English name of Buff-streaked Chat to Buff-streaked Bushchat as the word "Bushchat" is specific to the genus. Ibid.

July 15, 2003. Page 492

Gray-headed Robin (*Heteromyias cinereifrons*) is now considered a race of *Heteromyias albispectus*. Change English name of *H. albispectus* to Gray-headed Robin. Higgins, P. J. and J. M. Peter, editors. 2002. Handbook of Australian, New Zealand and Antarctic Birds. Volume 6: Pardalotes to shrike-thrushes. Melbourne: Oxford University Press.

July 15, 2003. Page 500

Add race *sweeti* to Black-hooded Laughingthrush. Range of nominate *G. m. milleti*: Montane forests of n and s Vietnam and s Laos; range of *G. m. sweeti*: Central Highlands of Vietnam (Kon Tum Province). Eames, Jonathan. 2002. Eleven new sub-species of babbler (*Passeriformes: Timaliinae*) from Kon Tum Province, Vietnam. Bulletin B. O. C. 122 (2): 109-141.

July 15, 2003. Page 507

Under Coral-billed Scimitar-Babbler add race *dickinsoni* after *P. f. orientalis*. Range: Highlands of central Vietnam and Bolavens Plateau of s Laos. Ibid.

July 15, 2003. Page 515

Under *Cutia* add race *hoae* after *C. n. cervinocrissa*. Range: Central Vietnam (Mt. Ngoc Linh, Kon Tum Province). Ibid.

July 15, 2003. Page 517

Under Chestnut-tailed Minla add race *traii* after *M. s. malayana*. Range: Central Vietnam (Mt. Ngoc Linh, Kon Tum Province). Ibid.

Under Golden-breasted Fulvetta add race *robsoni* after *A. c. amoena*. Range: Central Vietnam (Mt. Ngoc Linh, Kon Tum Province). Ibid.

Under Rufous-winged Fulvetta add race *stepanyani* after *A. c. soror*. Range: Highlands of c Vietnam (Kon Tum Province and Gia Lai Province). Ibid.

July 15, 2003. Page 518

Add race *cui* to Rusty-capped Fulvetta. Range of nominate *A. d. dubia*: Pine forests of s Myanmar and adj. China (Yunnan). Range of *A. d. cui*: Range: Central Vietnam (Mt. Ngoc Linh, Kon Tum Province). Ibid.

July 15, 2003. Page 519

Under Rufous-backed Sibia add race *roundi* after *H. a. mixta*. Range: Central Vietnam (Kon Tum Province). Ibid.

Under Black-backed Sibia add race *kingi* after *H. m. robinsoni*.

Range: Central Vietnam (Mt. Ngoc Linh, Kon Tum Province). Ibid.

July 15, 2003. Page 520

Under Stripe-throated Yuhina add race *uthaii* after *Y. g. gularis*.

Range: Central Vietnam (Mt. Ngoc Linh, Kon Tum Province). Ibid.

July 15, 2003. Page 522

Under Black-throated Parrotbill add race *kamoli* after *P. n. feae*.

Range: Central Vietnam (Mt. Ngoc Linh, Kon Tum Province). Ibid.

July 15, 2003. Page 523

Logrunner is split into two species. Northern Logrunner *Orthonyx novaeguineae* and Southern Logrunner *Orthonyx temminckii*. Higgins, P. J. and J. M. Peter, op. cit.

July 15, 2003. Page 523 and xvii

Change family name of Whipbirds and Quail-thrushes from Cinclosomatidae to Eupetidae. Eupetidae Bonaparte, 1850 is senior to Cinclosomatidae Mathews, 1921 as the family name for the whipbird—quail-thrush group (Bock 1994; pace Sibley & Ahlquist 1990; Sibley & Monroe 1990; Christidis and Boles 1994). Schodde, R. and I. J. Mason. 1999. The Directory of Australian Birds: Passerines: 407-408. Collingwood, Victoria: Csiro Publishing.

July 15, 2003. Page 528

Change English name of *Sericornis humilis* to Tasmanian Scrubwren.

Add race *S. h. tregallasi* with range King Island (Bass Strait).

Higgins, P. J. and J. M. Peter, op. cit.

July 15, 2003. Page 536

Race *guineensis* of White-winged Black Tit is raised to species level. White-shouldered Black Tit *Melaniparus guineensis*. Range as published in the checklist. Borrow, Nik and Ron Demey. 2001. A Guide to the Birds of Western Africa. Princeton University Press.

July 15, 2003. Page 539

The Blue Tit is split into two species: Eurasian Blue Tit *Cyanistes caeruleus* and African Blue Tit *Cyanistes teneriffae*. Following races belong to the *Cyanistes teneriffae* complex: *ultramarinus*, *cyrenaicae*, *palmensis*, *teneriffae*, *ombriosus*, and *degener*. Ranges as published in the checklist. Salzburger, Walter, Jochen Martens and Christian Sturmbauer. 2002. Paraphyly of the Blue Tit (*Parus*

caeruleus) suggested from cytochrome b sequences. Molecular Phylogenetics and Evolution Vol. 24 (1) 19-25.

July 15, 2003. Page 542

After Eurasian Treecreeper add Sichuan Treecreeper *Certhia tianquanensis*. Range: Known only from mountains of s China (Sichuan Province). Martens, J., S. Eck and Y-H Sun. *Certhia tianquanensis* Li, a treecreeper with relict distribution in Sichuan, China. 2002. Journal Ornithology 143: 440-456.

July 15, 2003. Page 552

Race *bella* of Lovely Sunbird *Aethopyga bella* is raised to species level with six subspecies. Race *shelleyi* becomes monotypic with English name Handsome Sunbird *Aethopyga shelleyi*. Mann, C. F. (in prep.). Systematic notes on Asian birds. Taxonomic comments on some south-east Asian members of the Nectariniidae. Zool. Verh. Leiden. Oriental Bird Club Bulletin 36: 15-16.

July 15, 2003. Page 604

The Tibetan Ground-Jay *Pseudopodoces humilis* is removed from the Corvidae and placed in the Paridae after the Sultan Tit on p. 539. Change the English name to Ground Tit. James, Helen F., Per G. P. Ericson, Beth Slikas, Fu-Min Lei, Frank B. Gill and Storrs L. Olsen. 2003. *Pseudopodoces humilis*, a misclassified terrestrial tit (Paridae) of the Tibetan Plateau: evolutionary consequences of shifting adaptive zones. Ibis 145 (2): 185-202.

July 15, 2003. Page 636

Change range of *Vireo altiloquus barbatulus* to read: Coastal s Florida, Cuba and Haiti; winters to Amazonian Brazil. There is no specimen evidence that *Vireo altiloquus* occurs in Peru. Schulenberg, Thomas S. and Paul Salaman. 2003. Specimen record of Black-whiskered *Vireo* *Vireo altiloquus* in Peru is erroneous. Bulletin B. O. C. 123 (1): 57-59.

July 15, 2003. Page 661

Race *auricularis* of Black-capped Hemispingus is raised to species level. White-browed Hemispingus *Hemispingus auricularis*. Range: Andes of e Peru (Amazonas to Cusco). Garcia-Moreno, J., J. Ohlson and J. Fjelds . 2001. MtDNA Sequences Support Monophyly of Hemispingus Tanagers. Molecular Phylogenetics and Evolution. Vol. 21 (3): 424-435.

Race *piurae* of Black-eared Hemispingus is raised to species level. Piura Hemispingus *Hemispingus piurae*. Range: Extreme sw Ecuador and nw Peru (Piura). Ibid.

July 15, 2003. Page 693

Race *arcticus* of Eastern Towhee is now considered a race of Spotted Towhee. Beadle, David and James Rising. 2002. Sparrows of the United States and Canada. Academic Press.

July 15, 2003. Page 696

Correct spelling of race *pelonotus* of Seaside Sparrow. David, Normand and Michel Gosselin. 2002. Gender agreement of avian species names. Bulletin B. O. C. 122: 14-49.

July 15, 2003. Page 697

Races *alter* and *subvirgatus* are considered races of Nelson's Sharp-tailed Sparrow. Ranges as published in the checklist. Beadle and Rising, op. cit. Note that correct spelling is *alter*, not *alterus*. David and Gosselin. Ibid.

Correct spelling of Henslow's Sparrow race *A. h. henslowii*.

July 15, 2003. Page 698

Correct spelling of Song Sparrow race *M. m. heermanni*.

Add race *M. g. nigrescens* under Swamp Sparrow. Range: Atlantic coast from New Jersey south to Maryland. Beadle and Rising, op. cit.

July 15, 2003. Page 699

Races *caniceps* and *dorsalis* listed under Yellow-eyed Junco are now considered races of Dark-eyed Junco *Junco hyemalis*. Beadle and Rising, op. cit.

July 15, 2003. Page 705

Change ranges of Tawny-shouldered Blackbird to read as follows: *A. h. humeralis*, Range: Cuba and Haiti; *A. h. scopulus*, Range: Cayo Cantilles (south of Cuba). Jaramillo, Alvaro and Peter Burke. 1999. New World Blackbirds: The Icterids. Princeton University Press.

July 15, 2003. Page 709

The Troupial is split into three species. Venezuelan Troupial *Icterus icterus* with races *ridgwayi*, *icterus* and *metae*. Ranges as listed in the checklist. Campo Troupial *Icterus jamaicaii*. Monotypic. Range: Lowlands of eastern Brazil. Orange-backed Troupial *Icterus croconotus* with races *croconotus* and *strictifrons*. Ranges as listed in the checklist. Jaramillo, Alvaro and Peter Burke. 1999. New World Blackbirds: The Icterids. Princeton University Press; and Omland, K.E., S. M. Lanyon and S. J. Fritz. 1999. A molecular phylogeny of the New World Orioles (*Icterus*): The importance of dense taxon sampling. Molecular Phylogenetics and Evolution. 12: 224-239.

July 15, 2003. Page 711

Change scientific name of the Bolivian Blackbird to *Agelaioides*

oreopsar. Lowther, Peter. 2001. New name for the Bolivian Blackbird. Bulletin B. O. C. 121 (4) 280.

December 15, 2003. Page 10

The race of the Herald Petrel breeding on Henderson Island is raised to species level. Henderson Petrel *Pterodroma atrata*. Brooke, M. de L. and G. Rowe. 1996. Behavioural and molecular evidence for the specific status of light and dark morphs of the Herald Petrel *Pterodroma heraldica*. Ibis 138 (3): 420-432.

December 15, 2003. Page 20

Delete Galapagos Heron. Now considered a race of widespread Striated Heron *Butorides striata**. Add under *B. s. striata*. Banks, Richard C., Carla Cicero, Jon L. Dunn, Andrew W. Kratter, Pamela C. Rasmussen, J. V. Remsen, Jr., James D. Rising and Douglas F. Stotz. 2003. Forty-fourth supplement to the American Ornithologists' Union Check-list of North American birds. The Auk 120 (3): 923-931.*Correct spelling is *striata*, not *striatus* contra AOU. "A compound genus-group name ending in the suffix...ides...is to be treated as masculine unless its author, when establishing the name, stated that it had another gender or treated it as such by combining it with an adjective species-group name in another gender form (ICZN 1999: Art. 30,.1.4.4). Therefore *Butorides* Blyth, 1852, was first established in combination with the feminine adjective *javanica*, and is thus feminine." David, Normand and Michel Gosselin. 2002. The grammatical gender of avian genera. Bull. B. O. C. 122 (4): 261.

December 15, 2003. Page 83

After Bare-eyed Rail add Talaud Rail *Gymnocrex talaudensis*. Range: Karekelong Island (Talaud Archipelago). Taylor, Barry. 1998. Rails: A guide to the Rails, Crakes, Gallinules and Coots of the World. Yale University Press.

December 15, 2003. Page 101

Change English name of *Larus belcheri* to Belcher's Gull. Banks, et al. 2003, op. cit.

December 15, 2003. Page 112-113

All New World pigeons in the genus *Columba* are changed to *Patagioenas*. The following species are affected: White-crowned Pigeon *Patagioenas leucocephala*; Scaly-naped Pigeon *Patagioenas squamosa*; Scaled Pigeon *Patagioenas speciosa*; Picazuro Pigeon *Patagioenas picazuro*; Bare-eyed Pigeon *Patagioenas corensis*; Spot-winged Pigeon *Patagioenas maculosa*; Band-tailed Pigeon *Patagioenas fasciata*; Chilean Pigeon *Patagioenas araucana*; Ring-tailed Pigeon *Patagioenas caribaea*; Pale-vented Pigeon *Patagioenas cayennensis*; Red-billed Pigeon *Patagioenas flavirostris*; Peruvian Pigeon *Patagioenas oenops*; Plain Pigeon *Patagioenas*

inornata; Plumbeous Pigeon *Patagioenas plumbea*; Short-billed Pigeon *Patagioenas nigrirostris*; Ruddy Pigeon *Patagioenas subvinacea* and Dusky Pigeon *Patagioenas goodsoni*. Banks, et al. 2003, op. cit.

December 15, 2003. Page 146

Due to a taxonomic revision the Painted Parakeet *Pyrrhura picta* is split into following species:

Azuero Parakeet *Pyrrhura eisenmanni*. Range: S-central Panama (Azuero Peninsula)

Sinu Parakeet *Pyrrhura subandina*. Range: NW Colombia (lower Sinú River Valley)

Todd's Parakeet *Pyrrhura caeruleiceps*. Range: W slope of Eastern Andes of n Colombia and Sierra de Perijá (Colombia/Venezuela border)

Painted Parakeet *Pyrrhura picta*:

P. p. picta. Range: SE Venezuela to the Guianas and n Amazonian Brazil (Amapá)

P. p. pantchenkoi. Range: Sierra de Perijá (Colombia/Venezuela border)

P. p. microtera. Range: N-central Brazil south of the Amazon (Pará to n Goiás)

Red-crowned Parakeet *Pyrrhura roseifrons*. Range: Disjunct in w Amazonia (w Brazil in Amazonas and n-central Peru s of Río Marañón to Huánuco)

Deville's Parakeet *Pyrrhura lucianii*. Range: Brazil (w-central Amazonia along Rio Solimões [at Tefe] and Rio Purús)

Hellmayr's Parakeet *Pyrrhura amazonum*. Range: N-central Brazil (e and se Amazonia in Pará, n Mato Grosso, n Goiás and w Maranhão)

Madeira Parakeet *Pyrrhura snethlage*. Range: Drainage of Rio Madeira in s Brazil and n Bolivia

Wavy-breasted Parakeet *Pyrrhura peruviana*. Range: Two disjunct populations, one in se Ecuador and n Peru and another in s-central Peru (Junín).

Joseph, L. 2000. Beginning an end to 63 years of uncertainty: The Neotropical parakeets known as *Pyrrhura picta* and *P. leucotis* comprise more than two species. *Proceedings of the Academy of Natural Sciences of Philadelphia* 150: 279-292; Joseph, L. and D. Stockwell. 2002. Climatic modeling of the distribution of some *Pyrrhura* parakeets of northwestern South America with notes on their systematics and special reference to *Pyrrhura caeruleiceps* Todd, 1947. *Ornitologia Neotropical* 13: 1-8; Joseph, L. 2002. Geographical variation, taxonomy and distribution of some Amazonian *Pyrrhura* parakeets. *Ornitologia Neotropical* 13: 337-363.

December 15, 2003. Page 166-168

All New World owls in the genus *Otus* are changed to the genus *Megascops*. Western Screech-Owl *Megascops kennicottii*; Balsas Screech-Owl *Megascops seductus*; Pacific Screech-Owl *Megascops cooperi*; Whiskered Screech-Owl *Megascops trichopsis*; Eastern

Screech-Owl *Megascops asio*; Tropical Screech-Owl *Megascops choliba*; Koepcke's Screech-Owl *Megascops koepckeae*; West Peruvian Screech-Owl *Megascops roboratus*; Bare-shanked Screech-Owl *Megascops clarkii*; Bearded Screech-Owl *Megascops barbarus*; Rufescent Screech-Owl *Megascops ingens*; Colombian Screech-Owl *Megascops colombianus*; Cinnamon Screech-Owl *Megascops petersoni*; Cloud-forest Screech-Owl *Megascops marshalli*; Tawny-bellied Screech-Owl *Megascops watsonii*; Guatemalan Screech-Owl *Megascops guatemalae*; Vermiculated Screech-Owl *Megascops vermiculatus*; Roraima Screech-Owl *Megascops roraimae*; Rio Napo Screech-Owl *Megascops napensis*; Hoy's Screech-Owl *Megascops hoyi*; Variable Screech-Owl *Megascops atricapilla*; Long-tufted Screech-Owl *Megascops sanctaecatarinae*; Puerto Rican Screech-Owl *Megascops nudipes*; White-throated Screech-Owl *Megascops albogularis*. Banks, et al. 2003, op. cit.

December 15, 2003. Page 168

Change English name of *Gymnoglaux lawrencii* to Bare-legged Owl. Banks, et al. 2003, op. cit.

December 15, 2003. Page 170

Change scientific name of Snowy Owl to *Bubo scandiacus* and move to p. 169 after Philippine Eagle-Owl. Banks, et al. 2003, op. cit.

December 15, 2003. Page 180

Change scientific name of Bahia Nighthawk to *Nyctiprogne vielliardi* and move to page 181 after Band-tailed Nighthawk. Add northern Minas Gerais to range. Whitney, Bret M., José F. Pacheco, Paulo Sérgio M. da Fonseca, Richard E. Webster, Guy M. Kirwan and Juan M. Barnett. 2003. Reassignment of *Chordeiles vielliardi* Lencioni-Neto, 1994, to *Nyctiprogne Bonaparte*, 1857, with comments on the latter genus and some presumably related *chordeilines* (Caprimulgidae). Bulletin B. O. C. 123 (2): 103-112.

December 15, 2003. Page 290

Change scientific name of Zimmer's Woodcreeper to *Dendroplex kienerii*. The name *necopinus* is now considered a junior synonym of *X. kienerii*. Change scientific name of Straight-billed Woodcreeper to *Dendroplex picus* and move both to follow Olive-backed Woodcreeper. Alexio, Alexandre. 2002. Molecular systematics and the role of the "Várzea"- "Terra-firme" ecotone in the diversification of *Xiphorhynchus* woodcreepers (Aves: Dendrocolaptidae). The Auk 119 (3): 621-640

Race *chunchotambo* of Ocellated Woodcreeper with race *brevirostris* raised to species level. Tschudi's Woodcreeper *Xiphorhynchus chunchotambo*. Range as published in the checklist. Ibid.

Spix's Woodcreeper is now considered monotypic. Race *elegans* with

subspecies *ornatus*, *buenavistae* and *insignis* is raised to species level. Elegant Woodcreeper *Xiphorhynchus elegans*. Ranges as published in the checklist. Range of race *buenavistae* is East slope of Eastern Andes of Colombia in Orinoco drainage. Ibid.

Race *juruanus* of Elegant Woodcreeper is raised to species status. Juruá Woodcreeper *Xiphorhynchus juruanus*. Range as published in the checklist. Raposo, Marcos A. and Elizabeth Höfling. 2003. Alpha taxonomy of the *Xiphorhynchus spixii* species group with the validation of *X. juruanus* Ihering, 1904. *Cotinga* 20: 72-80.

Race *guttatoides* of Buff-throated Woodcreeper is raised to species level. Lafresnaye's Woodcreeper *Xiphorhynchus guttatoides*, with following races: *eytoni*, *guttatoides*, *vicinialis* and *dorbignyanus*. Ranges as published in the checklist. Ibid.

December 15, 2003. Page 292

Move Lesser Woodcreeper to page 290 following Tschudi's Woodcreeper. Change genus from *Lepidocolaptes* to *Xiphorhynchus*. Ibid.

December 15, 2003. Page 292

Race *falcinellus* of Scaled Woodcreeper is raised to species level. Scalloped Woodcreeper *Lepidocolaptes falcinellus*. Range as published in the checklist. del Hoyo, J., A. Elliot and D. A. Christie, eds. 2003. *Handbook of the Birds of the World*. Vol. 8. Broadbills to Tapaculos. Barcelona: Lynx Edicions.

December 15, 2003. Page 405

After Gray-breasted Wood-Wren add: Munchique Wood-Wren *Henicorhina negreti*. Range: Western Andes of Colombia. Salaman, P., Paul Coopmans, T. M. Donegan, M. Mulligan, A. Cortés, S. L. Hilty and L. A. Ortega. 2003. *Ornitologia Colombiana* 1: 4-20.

December 15, 2003. Page 439

Race *tangorum* of Manchurian Reed-Warbler is now considered a race of Paddyfield Warbler *Acrocephalus agricola*. Range as published in the checklist.

December 15, 2003. Page 442

Change English name of *Stenostira scita* to Fairy Flycatcher and move to page 460 after Gray Tit-Flycatcher. Now regarded as a muscicapid flycatcher. Urban, E. K., C. H. Fry and S. Keith, editors. 1997. *The Birds of Africa*, Vol. 5. San Diego: Academic Press.

December 15, 2003. Page 452

The Long-legged Warbler *Trichocichla rufa*, has been rediscovered in Wabu Forest Reserve on Fiji. The news release from Fiji Birdlife was somewhat confusing since it states that the bird "was believed to

have been extinct for more than a century.” The subspecies *T. r. cluniei* was described in 1974 from the southern slopes of the Delanacau Mountains in west-central Vanua Levu. Kinsky, F. C. 1975. A new subspecies of the Long-legged Warbler *Trichocichla rufa* Reichenow, from Vanua Levu, Fiji. Bulletin B. O. C. 95: 98-101.

December 15, 2003. Page 460

Race *albicilla* of Red-breasted Flycatcher is raised to species level. Taiga Flycatcher *Ficedula albicilla*. Range as published in the checklist. Ibis 146: 153-156.

December 15, 2003. Page 461

Move Damar Flycatcher *Ficedula henrici* from page 714 to page 461 after Little Slaty Flycatcher. The bird was rediscovered in September 2001 after a 102-year absence on Damar Island in the eastern Lesser Sundas and found to be “widespread and common.” Shannaz, J. 2001. Rediscovery of the Damar Flycatcher *Ficedula henrici* on Damar Island, Maluku, Indonesia. Oriental Bird Club Bulletin 34: 38-39.

December 15, 2003. Page 617

Change English name of *Ploceus bertrandi* to Bertram’s Weaver. Dowsett, R. J. and J. A. Jobling. 2003. Bertram Lutley Sclater and Bertram’s Weaver *Ploceus bertrandi*. Bulletin B. O. C. 123 (2): 135-136.

December 15, 2003. Page 623

Move Parasitic Weaver to page 633 after Broad-tailed Paradise-Whydah. Lahti, David C. and Robert B. Payne. 2003. Morphological and behavioral evidence of relationships of the Cuckoo Finch *Anomalospiza imberbis*. Bulletin B. O. C. 123 (2): 113-125.

December 15, 2003. Page 641

Race *megaplaga* of White-winged Crossbill is raised to species level. Hispaniolan Crossbill *Loxia megaplaga*. Insert after Red Crossbill. Range as published in the checklist. Banks, et al. 2003, op. cit.

December 15, 2003. Page 642

Lesser Redpoll, which was split from the Common Redpoll in the December 31, 2000 update, is now lumped once again as the race *cabaret* of the Common Redpoll. Ottvall, R., S. Bensch, G. Walinder and J. T. Lifjeld. 2003. No evidence of genetic differentiation between Lesser Redpolls *Carduelis flammea cabaret* and Common Redpolls *Carduelis f. flammea*. Avian Science 2 (4): 237-244.

December 15, 2003. Page 685

After Blue Seed-eater add Carrizal Seed-eater *Amaurospiza carrizalensis*. Range: Known only from lower Río Caroni in Bolívar,

Venezuela. Lentino, Miguel and Robin Restall. 2003. The Auk 120 (3): 600-606. (Note: In keeping with the use of English names suggested by the late Burt Monroe, I have shortened the suggested name of Carrizal Blue-black Seedeater to the less-cumbersome Carrizal Seedeater).

December 15, 2003. Page 709

Race *fuertesi* of Orchard Oriole is raised to species level *Fuertes'* Oriole *Icterus fuertesi*. Race *phillipsi* is no longer considered a valid subspecies and becomes part of range of the monotypic *I. spurius*: Baker, Jason M., Esteban López-Medrano, Adolfo G. Navarro-Sigüenza, Octavio R. Rojas-Soto and Kevin E. Omland. 2003. Recent speciation in the Orchard Oriole group: divergence of *Icterus spurius spurius* and *Icterus spurius fuertesi*. The Auk 120 (3): 848-859.

July 15, 2004. Page 8

Delete Amsterdam Albatross. Better considered a race of Wandering Albatross *D. e. amsterdamensis*. Penhallurick, John and Michael Wink. Analysis of the taxonomy and nomenclature of the Procellariiformes based on complete nucleotide sequences of the mitochondrial cytochrome b gene. Emu 104: 125-147. *Despite the tendency to split many of the Diomedidae into as many as 26 species, under the Biological Species Concept we reject these recently proposed splits. I quote from Penhallurick and Wink's paper: "Somewhat surprising is the distance evidence relating to *D. amsterdamensis*, which has generally been treated as a good species since its description by Roux et al. (1983), although Bourne (1989)...treated it as a subspecies of *D. exulans*. The fact that it is only 0.52% (0.00%) distant from *antipodensis*, *gibsoni* and *exulans*, and only 0.87% (0.00%) removed from *dabbenena* strongly suggests that it belongs among the subspecies of *exulans*."

July 15, 2004. Page 8

Delete race *platei* under Buller's Albatross. Now considered monotypic. Race *platei* is known to be a juvenile plumage of *T. bulleri*. Ibid.

July 15, 2004. Page 9

Under Shy Albatross add race *T. c. steadi*. Range Auckland Islands. Change range of *T. c. cauta* to Tasmania and adjacent waters. Ibid.

July 15, 2004. Page 37

The Canary Islands form of the Egyptian Vulture is now considered a valid subspecies, *N. p. majorensis*. Donázar, J.A., J. J. Negro, C. J. Palacios, L. Gangoso, J. A. Godoy, O. Caballos, F. Hiraldo and N. Capote. 2002. Description of a new subspecies of the Egyptian Vulture (Acciptridae: *Neophron percnopterus*) from the Canary

Islands. Journal Raptor Research 36: 17-23.

July 15, 2004. Page 45

Variable Hawk is split into two species (contra Farquhar 1998).
Red-backed Hawk *Buteo polyosoma* (with race *exsul*) and Puna Hawk
Buteo poecilochrous. Ranges as published in the checklist. Cabot,
José and Tjitte de Vries. 2003. *Buteo polyosoma* and *Buteo*
poecilochrous are two distinct species. Bulletin B. O. C. 123 (3):
190-207.

July 15, 2004. Page 69

The rare Orange-necked Partridge, previously known from three
locations in south Vietnam, has been found in adjacent Cambodia.
Davidson, Peter. 2003. Endangered partridge discovered in Cambodia.
Oriental Bird Club Bulletin 37: 61.

July 15, 2004. Page 71

Under Sclater's Monal add subspecies *L. s. arunachalensis*. Range:
Mts. of western Arunachal Pradesh (n India). Kumar, Suresh R. and
Pratap Singh. 2004. A new subspecies of Sclater's Monal *Lophophorus*
sclateri from western Arunachal Pradesh, India. Bull. B. O. C. 124
(1): 16-28.

July 15, 2004. Page 110

English name of *Columba livia* is changed to Rock Pigeon. Correct
spelling of *Columba livia gymnocyclus*. Change English name of *Columba*
oenas to Stock Dove. Banks, R. C. et al. 2002. Forty-third
supplement to the American Ornithologists' Union Check-list of North
American Birds. The Auk 119 (3): 897-906.

July 15, 2004. Page 146

Remove race *pantchenkoi* from Painted Parakeet. This form has been
regarded as a race of *caeruleiceps* but is now believed to be a worn
example the latter. Joseph, L. 2002. Geographical variation,
taxonomy and distribution of some Amazonian *Pyrrhura* parakeets.
Ornitologia Neotropical 13: 337-363.

July 15, 2004. Page 152

Under Guinea Tauraco change race *T. p. phoebus* to read *T. p.*
zenkeri. del Hoyo et al. Handbook of Birds of the World, Vol. 4.

July 15, 2004. Page 164

After Reddish Scops-Owl add: Serendib Scops-Owl *Otus thilohoffmanni*.
Range: Lowland rain forests of southwest Sri Lanka. Warakagoda,
Deepal H. and Pamela Rasmussen. 2004. A new species of scops-owl
from Sri Lanka. Bulletin B. O. C. 124 (2): 85-105.

July 15, 2004. Page 181

Diabolical Nightjar, known from a single 1931 specimen from northeast Sulawesi, was tape recorded and photographed in Lore Lindu National Park in central Sulawesi. Mauro, Iwein. 2003. Little-known Oriental Bird: Heinrich's Nightjar *Caprimulgus diabolicus*. Oriental Bird Club Bulletin 37: 30-34.

July 15, 2004. Page 245

Under Andean Toucanet race *cyanolaemus* should be a race of Black-throated Toucanet *Aulacorhynchus atrogularis*. Navarro, Adolfo G., A. Townsend Peterson, Esteban López-Medrano and Hesequio Benítez-Díaz. 2001. Species limits in Mesoamerican *Aulacorhynchus* Toucanets. Wilson Bulletin 114 (4): 363-372

July 15, 2004. Page 298

Race *ignota* of Pygmy Antwren is raised to species level. Moustached Antwren *Myrmotherula ignota*. Delete Short-billed Antwren *M. obscura*; now considered a race of *M. ignota*. del Hoyo, J., A. Elliott and D. A. Christie, eds. 2003. Handbook of the Birds of the World. Vol. 8. Broadbills to Tapaculos. Barcelona: Lynx Edicions.

July 15, 2004. Page 308

Race *monileger* of Black-faced Antthrush with races *pallidus* and *intermedius* is raised to species level. Mexican Antthrush *Formicarius monileger*. del Hoyo, et al. 2003. op. cit.

July 15, 2004. Page 314

Delete Perija Tapaculo. Now considered a race of White-crowned Tapaculo. del Hoyo, et al. 2003. op. cit.

July 15, 2004. Page 314

Delete Colombian Tapaculo. Now considered a race of Matorral Tapaculo. Change range of Matorral Tapaculo to read: E Andes of c Colombia (Cundinamarca and Boyacá). del Hoyo, et al. 2003. op. cit.

July 15, 2004. Page 314

After Tschudi's Tapaculo add: Lara Tapaculo *Scytalopus fuscicauda*. Range: Andes of Venezuela in Lara and Trujillo. del Hoyo, et al. 2003. op. cit.

July 15, 2004. Page 465

After Iringa Akalat add: Rubeho Akalat *Sheppardia aurantiithorax*. Range: Montane forests of e Tanzania (Rubehos and Ukagura Mts.). Beresford, Pamela, Jon Fjeldsø and Jacob Kiure. 2004. A new species of akalat (*Sheppardia*) narrowly endemic in the eastern arc of Tanzania. TheAuk 121 (2): 23-34

July 15, 2004. Page 552

Correct the following typographic error that appeared in the July 15, 2003 update. The monotypic Lovely Sunbird becomes *Aethopyga shelleyi*. Race *bella* is raised to species level as Handsome Sunbird *Aethopyga bella* with subspecies *flavipectus*, *minuta*, *rubrinota*, *bella*, *bonita* and *arolasi*. Ranges as published in the checklist. Note that the English name Shelley's Sunbird is preoccupied by the East African *Cinnyris shelleyi*. Mann, C. F. Systematic notes on Asian birds. Taxonomic comments on some south-east Asian members of the Nectariniidae. Zool. Verh. Leiden. Oriental Bird Club Bulletin 36: 15-16.

July 15, 2004. Page 614

Under *Passer diffusus* add race *P. d. stygiceps*; correct spelling of *P. d. mosambicus*.

July 15, 2004. Page 614

Rufous Sparrow is split into four species: Great Rufous Sparrow *Passer motitensis* with races *benguellensis* and *subsolanus*; Kenya Rufous Sparrow *Passer rufocinctus*; Shelley's Rufous Sparrow *Passer shelleyi*; and Kordofan Rufous Sparrow *Passer cordofanicus*. Ranges as published in the checklist. Fry, C. H. and Keith, S. (Eds). 2004. The Birds of Africa, Vol. VII. Princeton University Press.

July 15, 2004. Page 614

Change English name of *Passer melanurus* to Southern Gray-headed Sparrow. Race *P. m. vicinus* now considered a valid subspecies. Fry and Keith, op. cit.

July 15, 2004. Page 615

Under Yellow-throated *Petronia* add following races and ranges.

P. s. rufitergum: S Zaire to sw Tanzania, n Malawi, Zambia, Angola, nw Botswana

P. s. flavigula: S Zambia, Zimbabwe, w Mozambique, e Botswana and n S Africa

P. s. bororensis: E Tanzania, s Malawi, n Mozambique to ne Natal and e Swaziland

P. s. superciliaris: South Africa (e Cape Province to Natal)

Under Bush *Petronia* delete race *buchani*. Now considered monotypic. Fry and Keith, op. cit.

July 15, 2004. Page 616

Under Red-billed Buffalo-Weaver race *militaris* now considered synonymous with *B. n. niger*. Fry and Keith, op. cit.

July 15, 2004. Page 618

Change English name of *Ploceus velatus* to Southern Masked-Weaver, with following races:

P. v. velatus: Western Cape, Northern Cape and Free State

P. v. nigrifrons: E Cape, n and w Natal and w Swaziland
P. v. tahatali: Zimbabwe, se Botswana, sw Mozambique, e Swaziland, ne Natal and Northern Province
P. v. shelleyi: Zambia, Malawi and Mozambique
P. v. caurinus: S Angola, Namibia, Botswana and n Cape Province
P. v. finschi: Coastal Angola
Fry and Keith, op. cit.

July 15, 2004. Page 618

Race *vitellinus* of African Masked-Weaver with race *uluensis* is raised to species level, Vitelline Masked-Weaver *Ploceus vitellinus*.
Fry and Keith, op. cit.

July 15, 2004. Page 618

Race *ruweti* of Lufira Masked-Weaver is raised to species level. Lake Lufira Weaver *Ploceus ruweti*. Fry and Keith, op. cit..

July 15, 2004. Page 618

Change English name of *Ploceus reichardi* to Tanganyika Masked-Weaver. Fry and Keith, op. cit.

July 15, 2004. Page 618

Race *katangae* of African Masker Weaver is raised to species level. Katanga Masked-Weaver *Ploceus katangae*. Fry and Keith, op. cit.

July 15, 2004. Page 620

Change English name of Yellow-legged Malimbe to Yellow-legged Weaver and change scientific name to *Ploceus flavipes*. Fry and Keith, op.

July 15, 2004. Page 623

Race *jamesoni* of *Parmoptila rubrifrons* is elevated to species status. Jameson's Antpecker *Parmoptila jamesoni*. Change English name of *P. rubrifrons* to Red-fronted Antpecker. Woodcock, Martin W. 2003. Systematics and confusion in the genus *Parmoptila*. Bull. B. O. C. 123 (4): 274-277

July 15, 2004. Page 624

Race *lineata* of Red-winged *Pytilia* is raised to species status. Red-billed *Pytilia* *Pytilia lineata*. Range: Highlands of Ethiopia and e Sudan. Fry and Keith, op. cit.

July 15, 2004. Page 627

Under Black-headed Waxbill change name of race *graueri* to *marunguensis*. Form *kandti* is raised to species level. Kandt's Waxbill *Estrilda kandti*. Range: Rift Valley Mts. of Uganda, Rwanda and Kenya. Fry and Keith, op. cit.

July 15, 2004. Page 627

Change scientific name of Zebra Waxbill to *Sporaeginthus subflavus*.
Fry and Keith, op. cit.

July 15, 2004. Page 628

Race *fuscocrissa* of African Quailfinch with races *muelleri*, *smithersi*, *digressa* and *pallida* is raised to species level. African Quailfinch *Ortygospiza fuscocrissa*. Change English name of *Ortygospiza atricollis* to Black-faced Quailfinch. Under Black-faced Quailfinch race *ugandae* now considered synonymous with nominate *atricollis*. Fry and Keith, op. cit.

July 15, 2004. Page 628

Change genus of Locustfinch to *Paludipasser*. Fry and Keith, op. cit.

July 15, 2004. Page 629

Change genus of African Silverbill to *Euodice*. Fry and Keith, op. cit.

July 15, 2004. Page 630

Change genus of Gray-headed Silverbill to *Odontospiza*. Change scientific name of Bronze Mannikin to *Spermestes cucullatus* with race *scutatus*.

Change genus of Black-and-white Mannikin to *Spermestes*.

Change genus of Magpie Mannikin to *Spermestes*. Fry and Keith, op. cit.

July 15, 2004. Page 638

Race *louisae* of Golden-winged Grosbeak is raised to species level. Somali Grosbeak *Rhynchostruthus louisae*. Mts. of n Somalia near Ethiopian border. Fry and Keith, op. cit.

July 15, 2004. Page 644

African Citril is split into three species. African Citril *Serinus citrinelloides* with race *kikuyensis*; Western Citril *Serinus frontalis*; and Southern Citril *Serinus hypostictus* with race *brittoni*. Delete race *martinsi*. Fry and Keith, op. cit.

July 15, 2004. Page 644

Race *flavivertex* of Cape Canary with races *sassii* and *huillensis* is raised to species status. Yellow-crowned Canary *Serinus flavivertex*. Range: Highlands of Eritrea, Ethiopia, Kenya, n Tanzania, w Angola. Fry and Keith, op. cit.

July 15, 2004. Page 645

Under Black-throated Canary add race *seshekeensis*. Range: SW Zimbabwe. Race *reichenowi* is raised to species level Reichenow's Seedeater *Serinus reichenowi*. Fry and Keith, op. cit.

July 15, 2004. Page 646

Race *whytii* of Streaky Seedeater is raised to species level.
Yellow-browed Seedeater *Serinus whytii*; race *affinis* now considered synonymous with nominate *striolatus*. Fry and Keith, op. cit.

July 15, 2004. Page 646

Race *melanochrous* of Thick-billed Seedeater is raised to species level. Tanzania Seedeater *Serinus melanochrous*. Range: S Tanzania (Mt. Rungwe and Udzungwa Mts.). Fry and Keith, op. cit.

July 15, 2004. Page 646

Change genus of Cape Siskin and Drakensberg Siskin to *Pseudochloroptila*. Fry and Keith, op. cit.

July 15, 2004. Page 646

Change genus of Black-headed Canary to *Alario*. Fry and Keith, op. cit.

July 15, 2004. Page 646

Race *leucolaemus* of Black-headed Canary is raised to species level. Damara Canary *Alario leucolaemus*. Fry and Keith, op. cit.

July 15, 2004. Page 646

Change genus of Ankober Serin to *Carduelis* and move to page 644 to follow Warsangli Linnet. Fry and Keith, op. cit.

July 15, 2004. Page 648

Change genus of Desert Finch to *Rhodospiza* and genus of Trumpeter Finch to *Bucanetes*. Fry and Keith, op. cit.

July 15, 2004. Page 678

Race *vincenti* of Cape Bunting is raised to species status. Vincent's Bunting *Emberiza vincenti*. Fry and Keith, op. cit.

July 15, 2004. Page 689-690

Because of a major taxonomic revision the species sequence and ranges between Rufous-naped Brush-Finch and Ochre-breasted Brush-Finch are changed as follows:

Rufous-naped Brush-Finch *Atlapetes latinuchus*

A. l. *elaeoprurus* Subtropical n Central Andes of Colombia (Antioquia)

A. l. *simplex* E Andes of Colombia (Bogotá)

A. l. *caucae* Western and Central Andes of Colombia (Valle and Cauca)

A. l. *spodionotus* Andes of s Colombia (Nariño) and n Ecuador

A. l. *comptus* Andes of sw Ecuador (Cañar to w Loja) and n Peru (Piura)

A. l. latinuchus Andes of se Ecuador (e Azuay and e Loja) to ne Peru (Amazonas)
A. l. chugurensis Pacific slope of nw Peru (Cajamarca)
A. l. baroni N Peru (upper Mara—n Valley in Cajamarca and La Libertad)
 Vilcabamba Brush-Finch *Atlapetes terborghi* Andes of se Peru (Cordillera Vilcabamba)
 Black-faced Brush-Finch *Atlapetes melanolaimus* Subtropical Andes of se Peru (Cuzco and Puno)
 Bolivian Brush-Finch *Atlapetes rufinucha*
A. r. rufinucha Subtropical Andes of w Bolivia (La Paz and Cochabamba)
A. r. carrikeri Andes of e Bolivia (Santa Cruz)
 Slaty Brush-Finch *Atlapetes schistaceus*
A. s. schistaceus W and Central Andes of Colombia and Ecuador
A. s. tamae Andes of e Colombia and sw Venezuela (Ṭchira)
A. s. fumidus Sierra de Perij̣ (Colombia/Venezuela border)
A. s. castaneifrons Andes of w Venezuela (Trujillo, Mérida and e Ṭchira)
A. s. taczanowskii Andes of central Peru (Hụnuco and Jun'n)
 Cusco Brush-Finch *Atlapetes canigenis* Andes of se Peru (Cuzco)
 White-rimmed Brush-Finch *Atlapetes leucopis* Locally in Andes of s Colombia and e Ecuador
 Rufous-capped Brush-Finch *Atlapetes pileatus*
A. p. dilutus Oak-pine forests of n Mexican plateau
A. p. pileatus Oak-pine forests of s Mexican plateau
 Santa Marta Brush-Finch *Atlapetes melanocephalus* Santa Marta Mountains (ne Colombia)
 Olive-headed Brush-Finch *Atlapetes flaviceps* Known from two specimens from n Colombia (last recorded 1967)
 Dusky-headed Brush-Finch *Atlapetes fuscoolivaceus* Andes of Colombia (upper Magdalena Valley)
 Tricolored Brush-Finch *Atlapetes tricolor*
A. t. crassus Andes of w Colombia and Ecuador (Pichincha and El Oro)
A. t. tricolor Andes of central Peru (La Libertad and Jun'n)
 Moustached Brush-Finch *Atlapetes albofrenatus*
A. a. albofrenatus E Andes of Colombia (Norte de Santander to Cundinamarca)
A. a. meridae Andes of w Venezuela (Ṭchira and Mérida)
 Bay-crowned Brush-Finch *Atlapetes seebohmi*
A. s. celicae Andes of extreme s Ecuador (w Loja)
A. s. simonsi Andes of extreme s Ecuador (eastern and central Loja)
A. s. seebohmi Pacific slope of Andes of nw Peru (La Libertad to Ancash)
 Rusty-bellied Brush-Finch *Atlapetes nationi*
A. n. nationi Western Andes of Peru (Ancash to Arequipa)
A. n. brunneiceps Andes of sw Peru (Ica and Ayacucho)

White-winged Brush-Finch *Atlapetes leucopterus*
 A. l. *leucopterus* Andes of w Ecuador (south to Chimbo Valley)
 A. l. *dresseri* Andes of sw Ecuador (El Oro and Loja)
 A. l. *paynteri* N Peru (n Piura to Cajamarca border)
 White-headed Brush-Finch *Atlapetes albiceps* Arid scrub of se
 Ecuador and nw Peru
 Pale-headed Brush-Finch *Atlapetes pallidiceps* Arid Andes of sw
 Ecuador (Azuay)
 Rufous-eared Brush-Finch *Atlapetes rufigenis* Andes of nw Peru
 (Cajamarca, La Libertad, Ancash and Huánuco)
 Apurímac Brush-Finch *Atlapetes forbesi* Andes of s-central Peru
 (Apurímac, Cuzco and Puno)
 Black-spectacled Brush-Finch *Atlapetes melanopsis* Andes of s Peru
 (Ayacucho and Huancavelica)
 Ochre-breasted Brush-Finch *Atlapetes semirufus*. Clements, James F.
 and Noam Shany. 2001. A Field Guide to the Birds of Peru. Temecula,
 CA: Ibis Publishing Company.